1

Table 1. Complete collected data on drug resistance.

Strain	Vancomycin	Meticillin	Tetracycline	Ciprofloxacin	Clindamycin	Erythromycin	Oxacillin	Gentamicin	Chloramphenicol	Kanamycin	Co-trimoxazole	Rifampicin	Daptomycin	Bacitracin	Penicillin	Spectinomycin
04-02981	S [57]	R [57]	S [57]	R [57]	R [57]	R [57]	R [57]	S [57]	R [57]						R [57]	
06BA18369		S [53]														
11819-97		R [79]														
16K									R [86]		S [86]	R [86]				
55/2053		S [69]														
58-424			R [60]			R [60]									R [60]	R [60]
65-1322			R [60]			R [60]									R [60]	R [60]
68-397			R [60]			R [60]									R [60]	R [60]
6850		S [24]														
A017934/97		R [69]														
A10102	S : 1 [64]												S : 0.5 [64]			
A5948	S : 1 [64]												S : 1 [64]			
A6224	S : 2 [71]															
A8115	S : 1 [64]												S : 0.5 [64]			
A8117	S : 1 [64]												[illegible]			
A8796	S : 1 [64]												[illegible]			
A8819	S : 1 [64]												[illegible]			
A9299	S : 1 − 2 [64]												[illegible]			
A9635	S : 1 [66]						S : 0.25 [66]						[illegible]			
A9719	S : 1 − 2 [64]												[illegible]			
A9754	S : 2 [64]												[illegible]			
A9763	S : 1 [64]												[illegible]			
A9765	S : 1 [64]												[illegible]			
A9781	S : 1 [64]												[illegible]			
Bmb9393		R [11]														
Btn1260		S [69]														
C101		S [69]														
CA-347		R [78]														
CBD-635	I [6]	R [6]	R [6]	R [6]	R [6]	R [6]		R [6]	R [6]							
CF-Marseille		R [70]	R [70]			R [70]		S [70]		R [70]						
CN1		R [8]			R [8]	R [8]										
[illegible]																
aureus VRS1	R [42]															
aureus VRS10	R [42]															
aureus VRS11a	R [42]															
aureus VRS11b	R [42]															
aureus VRS2	R [42]															
aureus VRS3a	R [42]															
aureus VRS4	R [42]															
aureus VRS5	R [42]															
aureus VRS6	R [42]															
aureus VRS7	R [42]															
aureus VRS8	R [42]															
aureus VRS9	R [42]															

Strain	Mupirocin	Teicoplanin	Fusidic acid	Trimethoprim	Moxifloxacin	Linezolid	Tigecycline	Fosfomycin
04-02981	S [57]	S [57]	S [57]		R [57]	S [57]	S [57]	S [57]
16K				S [86]				

Further table columns: Levofloxacin, Cefazolin, Tobramycin, Minocycline, Doxycycline, Neomycin, Cefotaxime, Sparfloxacin, Cephalexin, Clarithromycin, Ofloxacin, Nisin, Imipenem, Lincomycin, Quinupristin/dalfopristin, Ceftriaxone, Cholamphenicol, Ertapenem, Nourseothricin, Netilmicin, Lactoferricin, Tosufloxacin, Centhromycin, Chloramphenicol B, Norfloxacin, Ceftizoxime, Streptomycin, Temafloxacin, Nitrofurantoin, Cefoxitin, Lomefloxacin, Pazufloxacin, Sulfamethoxazole, Gramicidin, Cephalothin, Ampicillin, IB-367, Fleroxacin, Cotrimoxazole, Novobiocin, Azithromycin, Sisomicin, Clofazimine, Amikacin, Flucloxacillin, Enoxacin, Polymyxin B — [illegible]

Comments

- legend: red: drug resistant; green: drug susceptible; yellow: intermediate drug resistant; gray: contrary information of no CLSI interpretation; white: not information is found,
- drug susceptibility information for the strain TW20 was confirmed by e-mail correspondence with the authors of [20]
- drug susceptibility information for the strain ST398 was confirmed by e-mail correspondence with the authors of [74]
- accordingly to [31] 'ET3-1 (RF122) is susceptible to most antibiotics, excluding Spectinomycin', according to [56] JH1 is susceptible to Rifampicin, with MIC 0.012 $\mu g/ml$. We use information from [56], treating it as more reliable, because this study is newer and provides the exact MIC value.

References

1. Tadashi Baba, Taeok Bae, Olaf Schneewind, Fumihiko Takeuchi, and Keiichi Hiramatsu. Genome sequence of staphylococcus aureus strain newman and comparative analysis of staphylococcal genomes: Polymorphism and evolution of two major pathogenicity islands. *Journal of Bacteriology*, 190(1):300–310, January 2008.
2. Amanda M. Beltramini, Chitrangada D. Mukhopadhyay, and Vijay Pancholi. Modulation of cell wall structure and antimicrobial susceptibility by a staphylococcus aureus eukaryote-like Serine/Threonine kinase and phosphatase. *Infection and Immunity*, 77(4):1406–1416, April 2009. PMID: 19188361 PMCID: 2663143.
3. Tamar Bikels-Goshen, Elad Landau, Sam Saguy, and Roni Shapira. Staphylococcal strains adapted to epigallocathechin gallate (EGCG) show reduced susceptibility to vancomycin, oxacillin and ampicillin, increased heat tolerance, and altered cell morphology. *International journal of food microbiology*, 138(1-2):26–31, March 2010. PMID: 20132996.
4. Gilles Brackman, Paul Cos, Louis Maes, Hans J. Nelis, and Tom Coenye. Quorum sensing inhibitors increase the susceptibility of bacterial biofilms to antibiotics in vitro and in vivo. *Antimicrobial Agents and Chemotherapy*, 55(6):26552661, 2011.
5. Kellie Burnside, Annalisa Lembo, Melissa de los Reyes, Anton Iliuk, Nguyen-Thao BinhTran, James E. Connelly, Wan-Jung Lin, Byron Z. Schmidt, Anthony R. Richardson, Ferric C. Fang, Weiguo Andy Tao, and Lakshmi Rajagopal. Regulation of hemolysin expression and virulence of staphylococcus aureus by a Serine/Threonine kinase and phosphatase. *PLoS ONE*, 5(6):e11071, 2010.
6. Ronan K. Carroll, Whittney N. Burda, Jill C. Roberts, Kealy K. Peak, Andrew C. Cannons, and Lindsey N. Shaw. Draft genome sequence of strain CBD-635, a methicillin-resistant staphylococcus aureus USA100 isolate. *Genome Announcements*, 1(4):e00491–13, August 2013. PMID: 23846279.
7. Chih-Jung Chen, Clemens Unger, Wolfgang Hoffmann, Jodi A. Lindsay, Yhu-Chering Huang, and Friedrich Gtz. Characterization and comparison of 2 distinct epidemic community-associated methicillin-resistant staphylococcus aureus clones of ST59 lineage. *PLoS ONE*, 8(9):e63210, September 2013.
8. Feng-Jui Chen, Tsai-Ling Lauderdale, Lih-Shinn Wang, and I-Wen Huang. Complete genome sequence of staphylococcus aureus z172, a vancomycin-intermediate and daptomycin-nonsusceptible methicillin-resistant strain isolated in taiwan. *Genome announcements*, 1(6), 2013. PMID: 24309740 PMCID: PMC3853063.

[illegible]